arXiv:1702.01002v2 [math.CV] 10 Feb 2017

TOTALLY REAL EMBEDDINGS WITH PRESCRIBED POLYNOMIAL HULLS.

LEANDRO AROSIO[†] AND ERLEND FORNÆSS WOLD[††]

Abstract. We embed compact C^∞ manifolds into $\mathbb{C}^n$ as totally real manifolds with prescribed polynomial hulls. As a consequence we show that any compact C^∞ manifold of dimension d admits a totally real embedding into $\mathbb{C}^{\lfloor \frac{3d}{2} \rfloor}$ with non-trivial polynomial hull without complex structure.

1. Introduction

We will prove the following result.

Theorem 1.1. *Let $X \subset \mathbb{C}^m$ be a totally real C^∞ submanifold of dimension $d > 1$, and let $K \subset X$ be a compact subset. Let M be a compact C^∞ manifold of dimension d (possibly with boundary) and assume that there exists a C^∞ embedding $\phi : X \to M$. Then there exists a totally real C^∞ embedding $\psi : M \to \mathbb{C}^{m+\ell}$, where $\ell = \max\{\lfloor \frac{3d}{2} \rfloor - m, 0\}$, such that*

(i) $K \subset \psi(M)$,
(ii) $\widehat{\psi(M)} = \psi(M) \cup \widehat{K}$.

Our motivation for proving this result is the following recent result by Izzo and Stout.

Theorem 1.2 (Izzo–Stout [8])**.** *Let M be a compact C^∞ surface (possibly with boundary). Then M embeds into $\mathbb{C}^3$ as a totally real C^∞ surface Σ such that $\widehat{\Sigma} \setminus \Sigma$ is not empty and does not contain any analytic disc.*

In the case were M is a torus, Theorem 1.2 was proved by Izzo, Samuelsson and the second named author [7]. In a recent paper Gupta [5] gave an explicit example of an isotropic torus in $\mathbb{C}^3$ whose polynomial hull consists only of an annulus attached to the torus. We note that the torus embedded in [7] is also isotropic, the image being the graph of a real valued function on the torus in $\mathbb{C}^2$.

As a corollary to Theorem 1.1 we prove the following generalization of Theorem 1.2 to every dimension $d > 1$.

Date: February 13, 2017.
2010 *Mathematics Subject Classification.* 32E20.
Key words and phrases. Polynomial Convexity, Totally Real Manifolds.
[††]Supported by NRC grant no. 240569 .
[†]Supported by the SIR grant "NEWHOLITE - New methods in holomorphic iteration" no. RBSI14CFME.
Part of this work was done during the international research program "Several Complex Variables and Complex Dynamics" at the Center for Advanced Study at the Academy of Science and Letters in Oslo during the academic year 2016/2017.

Corollary 1.3. *Let M be a compact C^∞ manifold (possibly with boundary) of dimension $d > 1$. Then M embeds into $\mathbb{C}^{\lfloor \frac{3d}{2} \rfloor}$ as a totally real C^∞ submanifold Σ such that $\widehat{\Sigma} \setminus \Sigma$ is not empty and does not contain any analytic disc.*

A similar result was proved in [7] for dimension $2d + 4$ of the target space. Notice that by [6, Theorem 2.1] the dimension $\lfloor \frac{3d}{2} \rfloor$ is optimal for totally real embeddings of d-dimensional manifolds, which implies that Corollary 1.3 is optimal.

We emphasize that Theorem 1.1 is a simple consequence of the general fact that any C^∞ manifold of dimension d embeds into $\mathbb{C}^{\lfloor \frac{3d}{2} \rfloor}$ as a totally real submanifold, and the following general genericity result regarding polynomial hulls and totally real embeddings (to be proved in Section 4):

Theorem 1.4. *Let M be a compact C^∞ manifold (possibly with boundary) of dimension $d < n$ and let $f \colon M \to \mathbb{C}^n$ be a totally real C^∞ embedding. Let $K \subset \mathbb{C}^n$ be a compact polynomially convex set. Then for all $k \geq 1$ and for all $\varepsilon > 0$ there exists a totally real C^∞ embedding $f_\varepsilon \colon M \to \mathbb{C}^n$ such that*

1. *$\|f_\varepsilon - f\|_{C^k(M)} < \varepsilon$,*
2. *$f_\varepsilon - f = 0$ near $f^{-1}(K)$, and*
3. *$K \cup \widehat{f_\varepsilon(M)} = K \cup f_\varepsilon(M)$.*

In the case $K = \varnothing$, Theorem 1.4 was proved by Forstneric and Rosay for $d = 2$ and $n \geq 3$, by Forstneric [3] for $d \leq \frac{2n}{3}$, and by Løw and the second named author [9] for $d < n$.

To obtain Corollary 1.3 it is crucial that Alexander [2] has constructed a suitable compact set K in the totally real distinguished boundary of the bidisk in $\mathbb{C}^2$ (see Section 2).

2. Proof of Theorem 1.1 and Corollary 1.3

In this section we prove Theorem 1.1 and Corollary 1.3 modulo Theorem 1.4 which will be proved in the next section.

We recall first the following result [4, Lemma 5.3]. By *generic* we mean open and dense in the Whitney C^∞-topology.

Lemma 2.1. *Let M be a compact C^∞ manifold (possibly with boundary) of dimension d. Then a generic C^∞ embedding $\psi : M \to \mathbb{C}^{\lfloor \frac{3d}{2} \rfloor}$ is totally real.*

Proof of Theorem 1.1:

Let $U \subset\subset \phi(X) \subset M$ be an open subset with $\phi(K) \subset U$ and let $\psi_1 \colon \phi(X) \to \mathbb{C}^m$ be the inverse of ϕ. Let $\psi_2 \colon M \to \mathbb{C}^m$ be a C^∞ mapping which agrees with ψ_1 on U. Notice that since $2(m + \ell) \geq 2d + 1$, by Whitney's theorem [10, Theorem 2] a generic C^∞ mapping from M to $\mathbb{C}^{m+\ell}$ is an embedding, and since $m + \ell \geq \lfloor \frac{3d}{2} \rfloor$, by Lemma 2.1 a generic C^∞ embedding of M into $\mathbb{C}^{m+\ell}$ is totally real. Let $\psi_3 \colon M \to \mathbb{C}^{m+\ell}$ be a small perturbation of ψ_2 which is a totally real C^∞ embedding.

Let $\chi \in C_0^\infty(U)$ with $\chi \equiv 1$ near $\phi(K)$ and $0 \leq \chi \leq 1$. Then if the perturbation ψ_3 was small enough, the mapping from M to $\mathbb{C}^{m+\ell}$ defined as $\psi_4 := \chi \cdot \psi_1 + (1 - \chi)\psi_3$ is a totally real C^∞ immersion. Clearly $K \subset \psi_4(M)$. Since ψ_4 is injective in U, there exists a small C^∞ perturbation

$\psi\colon M \to \mathbb{C}^{m+\ell}$ which is a totally real C^∞ embedding and such that $\psi|_{\phi(K)} = \psi_4|_{\phi(K)}$ (see e.g. [1, Lemma 2.5]), which implies that $K \subset \psi(M)$. Since $m + \ell > d$, we can apply Theorem 1.4 with K replaced by $\widehat{K}$ and Theorem 1.1 follows.

□

To prove Corollary 1.3 we need the following result of Alexander:

Theorem 2.2. *(Alexander, [2]) The standard torus $\mathbb{T}^2 := \{(e^{i\vartheta}, e^{i\psi})\colon \vartheta, \psi \in \mathbb{R}\}$ in $\mathbb{C}^2$ contains a compact subset K such that $\widehat{K} \setminus K$ is not empty but contains no analytic discs. Such a set can be found in every neighborhood of the diagonal in $\mathbb{T}^2$.*

Proof of Corollary 1.3: Choose a small annular neighbourhood A of the diagonal in $\mathbb{T}^2$, and define a totally real C^∞ submanifold X of $\mathbb{C}^d$ by

$$X = \{(z_1, z_2, z_3, ..., z_d) : (z_1, z_2) \in A, \mathrm{Im}(z_j) = 0 \text{ and } -\epsilon < \mathrm{Re}(z_j) < \epsilon \text{ for } j = 3, ..., d\}.$$

Now X embeds into the into (any small portion) of the interior of M and A contains an Alexander set K. By Theorem 1.1, there exists a totally real C^∞ embedding $\psi\colon M \to \mathbb{C}^{\lfloor \frac{3d}{2} \rfloor}$ such that $K \subset \psi(M)$ and $\widehat{\psi(M)} = \psi(M) \cup \widehat{K}$. Since the hull $\widehat{K}$ cannot be contained in the totally real submanifold $\psi(M)$, the result follows from Theorem 2.2.

□

3. Proof of the main lemma

For any compact set $K \subset \mathbb{C}^n$ denote

$$h(K) := \overline{\widehat{K} \setminus K}.$$

The proof of the following main Lemma is essentially contained in [9, Proposition 6,7] where it is given in the case of C^1-regularity. We give a proof for completeness.

Lemma 3.1. *Let M be a compact C^∞ manifold (possibly with boundary) of dimension $d < n$ and let $f\colon M \to \mathbb{C}^n$ be a totally real C^∞ embedding. Let $K \subset \mathbb{C}^n$ be a compact polynomially convex set, and let U be a neighborhood of K. Then for all $k \geq 1$ and for all $\varepsilon > 0$ there exists a totally real C^∞ embedding $f_\varepsilon\colon M \to \mathbb{C}^n$ such that*

1. *$\|f_\varepsilon - f\|_{C^k(M)} < \varepsilon$,*
2. *$f_\varepsilon - f = 0$ near $f^{-1}(K)$, and*
3. *$K \cup \widehat{f_\varepsilon(M)} \subset U \cup f_\varepsilon(M)$.*

The following result is proved in [9, Proposition 4]).

Proposition 3.2. *Let M be a compact C^∞ manifold (possibly with boundary) and let $f\colon M \to \mathbb{C}^n$ be a totally real C^∞ embedding. Let $U' \subset\subset U \subset\subset \mathbb{C}^n$ be open sets. Then there exists an open neighborhood Ω of $f(M)$ such that*

1. *if $S \subset M$ is closed and $K \subset U'$ is compact, then*

$$h(K \cup f(S)) \subset U' \cup \Omega \Rightarrow h(K \cup f(S)) \subset U,$$

(2) *if f' is a sufficiently small C^1-perturbation of f, then (1) holds with S' in place of S for all closed subsets S' of M.*

The following lemma is proved in [9, Corollary 2]

Lemma 3.3. *Let $K \subset \mathbb{C}^n$ be a compact subset. Let M be a compact C^1 manifold (possibly with boundary) and let $f\colon M \to \mathbb{C}^n$ be a totally real C^1 embedding. Let $S \subset M$ be a closed subset. Let U be an open set such that $\widehat{K \cup f(S)} \subset U \cup f(S)$. Then there exists a constant $c > 0$ such that if $\|f - f'\|_{C^1} < c$, then $\widehat{K \cup f'(S)} \subset U \cup f'(S)$.*

Proof of Lemma 3.1: The proof is in two steps. Let I denote the interval $[0,1] \subset \mathbb{R}$.

First Step: The first step is proving the lemma in the case $M = I^d$. The proof is by induction on d. Fix a strictly positive $d \in \mathbb{N}$, and assume that the result holds for embeddings of I^{d-1}.

Let $U' \subset \mathbb{C}^n$ be an open set such that $K \subset U' \subset\subset U$. Let $K' \subset U'$ be a polynomially convex set such that $K \subset \operatorname{int}(K')$. Let us fix some notation.

(1) for $J \in \mathbb{N}$ and $\alpha \in \mathbb{N}^d$, $0 \le \alpha_j \le j-1$, we denote by I^J_α the cube
$$I^J_\alpha := \left[\frac{\alpha_1}{J}, \frac{\alpha_1+1}{J}\right] \times \cdots \times \left[\frac{\alpha_d}{J}, \frac{\alpha_d+1}{J}\right],$$
(2) for $1 \le m \le d$, $0 \le j \le J$, we denote by $G^J_{m,j}$ the $(d-1)$-dimensional cube
$$G^J_{m,j} := I \times \cdots \times I \times \underbrace{\left\{\frac{j}{J}\right\}}_{m\text{-th index}} \times I \cdots \times I,$$
(3) we denote by G^J the grid which is the union of all $(d-1)$-dimensional cubes
$$G^J := \bigcup_{m,j} G^J_{m,j},$$
(4) we denote by $G^J(\delta)$ the closed δ-neighborhood of the grid G^J,
$$G^J(\delta) := \{x \in I^d \colon \operatorname{dist}(x, G^J) \le \delta\},$$
(5) we denote by $Q^J_\alpha(\delta) \subset I^J_\alpha$ the smaller d-dimensional cube
$$Q^J_\alpha(\delta) := \overline{I^J_\alpha \setminus G^J(\delta)},$$
(6) we denote by S^J an n-tuple of distinct cubes I^J_α.

We will prove the following statement. There exists a big enough J, a small enough δ and a totally real C^∞ embedding $f_\varepsilon\colon M \to \mathbb{C}^n$ such that

(i) $\|f_\varepsilon - f\|_{C^k(I^d)} < \varepsilon$,
(ii) $f_\varepsilon - f = 0$ near $f^{-1}(K)$, and
(iii) $f_\varepsilon(I^J_\alpha) \cap K \neq \varnothing \Rightarrow f_\varepsilon(I^J_\alpha) \subset K'$,
(iv) $h(K' \cup f_\varepsilon(S^J) \cup f_\varepsilon(G^J(\delta))) \subset U$ for any choice of n-tuple S^J,
(v) for all α such that $f_\varepsilon(Q^J_\alpha(\delta)) \subset \mathbb{C}^n \setminus K$, there exists an entire function $g_\alpha \in \mathcal{O}(\mathbb{C}^n)$ such that $f_\varepsilon(Q^J_\alpha(\delta)) \subset \{g_\alpha = 0\} := Z(g_\alpha)$,
(vi) for all choices of $n+1$ different multi-indices $\alpha^1, \dots \alpha^{n+1}$ we have $\bigcap_{1 \le j \le n+1} Z(g_{\alpha^j}) = \varnothing$.

Once this is proved, the first step of the proof goes as follows. Assume that

$$q \in [K \cup f_\varepsilon(I^d)\widehat{]}.$$

Let μ be a representative Jensen measure for the linear functional on $\mathcal{P}(K \cup f_\varepsilon(I^d))$ defined as the evaluation at q, such that for all entire functions $g \in \mathcal{O}(\mathbb{C}^n)$ we have

$$\log|g(q))| \leq \int \log|g| \mathrm{d}\mu. \tag{3.1}$$

Assume that there exists α such that $\mu(f(Q^J_\alpha(\delta))) > 0$ and $f(Q^J_\alpha(\delta)) \cap K = \varnothing$. By (v) we get that $\int \log|g_\alpha| \mathrm{d}\mu = -\infty$, and by (3.1) we have that $q \in Z(g_\alpha)$. By (vi), there exists an n-tuple of cubes S^J such that the measure μ is concentrated on the subset

$$K' \cup f_\varepsilon(S^J) \cup f_\varepsilon(G^J(\delta)).$$

It follows that $q \in [K' \cup f_\varepsilon(S^j) \cup f_\varepsilon(G^J(\delta))\widehat{]}$, and by (iv) we obtain that $q \in U \cup f_\varepsilon(I^d)$.
Claim 1: If J is big enough then

$$h(K' \cup f(S^J)) \subset U' \tag{3.2}$$

for any choice of n-tuple S^J. Moreover, for a fixed big enough J, if f_0 is a sufficiently small C^1-perturbation of f, then (3.2) holds with f_0 in place of f.
Proof of Claim 1: Let U'' be an open neighborhood of K' with $U'' \subset\subset U'$. Let Ω' be given by Proposition 3.2 with data (f, U', U''). Assume by contradiction that there is a sequence $(J_t)_{t\in\mathbb{N}}$, $J_t \to \infty$ and a sequence S^{J_t} of n-tuples such that

$$h(K' \cup f(S^{J_t})) \not\subset U'.$$

Let $I^{J_t}_{\alpha^1(t)}, \ldots, I^{J_t}_{\alpha^n(t)}$ be the cubes in S^{J_t}. Up to passing to a subsequence we may assume that for any $1 \leq j \leq n$, the image $f(I^{J_t}_{\alpha^j(t)})$ converges to a point $q_j \in M$ as $t \to \infty$. Since $K' \cup \{q_j\}_{1\leq j\leq n}$ is polynomially convex, there exists a Runge and Stein neighborhood V of $K' \cup \{q_j\}_{1\leq j\leq n}$ such that $V \subset U'' \cup \Omega'$. For large enough t we have that $K' \cup f(S^{J_t}) \subset V$, which implies $h(K' \cup f(S^{J_t})) \subset U'' \cup \Omega'$, and thus by Proposition 3.2 we have a contradiction.

Since there are only a finite number of n-tuples S^J for a fixed J, the perturbation claim follows from Lemma 3.3. *End proof of Claim 1.*

Choose J big enough to ensure that Claim 1 holds and that for all α,

$$f(I^J_\alpha) \cap K \neq \varnothing \Rightarrow f(I^J_\alpha) \subset \mathrm{int}(K').$$

Claim 2: There exists a totally real C^∞ embedding $f_{\frac{\varepsilon}{2}} \colon I^d \to \mathbb{C}^n$ such that

(1) $\|f_{\frac{\varepsilon}{2}} - f\|_{C^k(I^d)} < \frac{\varepsilon}{2}$,
(2) $f_{\frac{\varepsilon}{2}} = f$ near $f^{-1}(K')$,
(3) $h(K' \cup f_{\frac{\varepsilon}{2}}(S^J) \cup f_{\frac{\varepsilon}{2}}(G^J)) \subset U'$ for any choice of n-tuple S^J.

Moreover, if f_0 is a sufficiently small C^1-perturbation of $f_{\frac{\varepsilon}{2}}$, then (3) holds with f_0 in place of $f_{\frac{\varepsilon}{2}}$.
Proof of Claim 2: Fix an n-tuple S^J. Choose an ordering of the $(d-1)$-dimensional cubes $(G^J_{m,j})_{m,j}$ and denote them by $G_1, \ldots, G_\ell$.

Fix $\eta > 0$. We will construct a family of totally real C^∞ embeddings $(f_j \colon I^d \to \mathbb{C}^n)_{0\le j\le \ell}$ such that

(a) $\|f_j - f_{j-1}\|_{C^k(I^d)} \le \eta$,
(b) $f_j = f_{j-1}$ near $f^{-1}(K') \cup S^J \cup \bigcup_{1\le i\le j-1} G_i$,
(c) $h\left(K' \cup f_j(S^J) \cup \bigcup_{1\le i\le j} f_j(G_i)\right) \subset U'$.

If we set $f_0 := f$, then (c) is true by Claim 1. Assume we constructed f_j, where $0 \le j \le \ell - 1$. Then by (c) the set $h(K' \cup f_j(S^J) \cup \bigcup_{0\le i\le j} f_j(G_i))$ is compact in U', so there exists a set U_1'' such that $K' \subset U_1'' \subset\subset U'$ satisfying $h(K' \cup f_j(S^J) \cup \bigcup_{1\le i\le j} f_j(G_i)) \subset U_1''$. Let Ω_1 be given by Proposition 3.2 with data (f_j, U', U_1''). Consider the totally real C^∞ embedding $f_j|_{G_{j+1}} \colon G_{j+1} \to \mathbb{C}^n$. Since

$$[K' \cup f_j(S^J) \cup \bigcup_{0\le i\le j} f_j(G_i)\widehat{]} \subset U_1'' \cup \Omega_1,$$

by the inductive assumption (the result holds for embeddings of I^{d-1}) there exists a totally real C^∞ embedding $\tilde{f}_j|_{G_{j+1}} \colon G_{j+1} \to \mathbb{C}^n$ such that

(1) $\|\tilde{f}_j - f_j\|_{C^k(G_{j+1})}$ is small,
(2) $\tilde{f}_j = f_j$ near $f_j^{-1}(K') \cup S^J \cup \bigcup_{1\le i\le j} G_i$,
(3) $h\left(K' \cup f_j(S^J) \cup \bigcup_{1\le i\le j} f_j(G_i) \cup \tilde{f}_j(G_{j+1})\right) \subset U_1'' \cup \Omega_1$.

If $\|\tilde{f}_j - f_j\|_{C^k(G_{j+1})}$ is small enough, then we may extend $\tilde{f}_j$ to a totally real C^∞ embedding $f_{j+1} \colon I^d \to \mathbb{C}^n$ such that $\|f_{j+1} - f_j\|_{C^k(I^d)} \le \eta$, which coincides with f_j on $f^{-1}(K') \cup S^J \cup \bigcup_{1\le i\le j} G_i$. By the choice of Ω_1 it follows that

$$h(K' \cup f_{j+1}(S^J) \cup \bigcup_{1\le i\le j+1} f_{j+1}(G_i)) \subset U'.$$

The mapping f_ℓ satisfies Claim 2 for the chosen n-tuple S^J.

We then iterate this argument for every n-tuple S^J (choosing η small enough), and we obtain $f_{\frac{\varepsilon}{2}}$. The perturbation claim follows from Lemma 3.3.
End proof of Claim 2.

Let Ω be an open neighborhood of $f_{\frac{\varepsilon}{2}}(I^d)$ given by Proposition 3.2 with data $(f_{\frac{\varepsilon}{2}}, U, U')$.
Claim 3: There exists a $\delta > 0$ such that

$$h(K' \cup f_{\frac{\varepsilon}{2}}(S^J) \cup f_{\frac{\varepsilon}{2}}(G^J(\delta))) \subset U \tag{3.3}$$

for any choice of n-tuple S^J. Moreover, if f_0 is a sufficiently small C^1-perturbation of $f_{\frac{\varepsilon}{2}}$, then (3.3) holds with f_0 in place of $f_{\frac{\varepsilon}{2}}$.
Proof of Claim 3: Fix an n-tuple S^J. Since $h(K' \cup f_{\frac{\varepsilon}{2}}(S^J) \cup f_{\frac{\varepsilon}{2}}(G^J)) \subset U'$, there exists a Runge and Stein neighborhood $V \subset\subset \Omega \cup U'$ of $K' \cup f_{\frac{\varepsilon}{2}}(S^J) \cup f_{\frac{\varepsilon}{2}}(G^J)$. If $\delta > 0$ is small enough we have that $K' \cup f_{\frac{\varepsilon}{2}}(S^J) \cup f_{\frac{\varepsilon}{2}}(G^J(\delta)) \subset\subset V$, and thus $[h(K' \cup f_{\frac{\varepsilon}{2}}(S^J) \cup f_{\frac{\varepsilon}{2}}(G^J(\delta)))\widehat{]} \subset U$. Since the number of n-tuples is finite, we can choose a δ which works for all of them. The perturbation claim follows from Lemma 3.3.

End proof of Claim 3.

We end Step 1 pushing each $f_{\frac{\epsilon}{2}}(Q_\alpha^J)$ with $f_{\frac{\epsilon}{2}}(Q_\alpha^J) \cap K = \varnothing$ into an analytic hypersurface of $\mathbb{C}^n$.

We will identify the cube I_α^J with a the cube centred at the origin in $\mathbb{R}^n$, with the decomposition $\mathbb{R}^d \oplus i\mathbb{R}^d \oplus \mathbb{C}^{n-d}$ of $\mathbb{C}^n$, and after a affine linear change of coordinates we may assume that $f_{\frac{\epsilon}{2}} : I_\alpha^J \to \mathbb{C}^n$ satisfies $f_{\frac{\epsilon}{2}}(0) = 0$, with the tangent space of $f_{\frac{\epsilon}{2}}(I_\alpha^J)$ at the origin equal to $\mathbb{R}^d$, and with $Df_{\frac{\epsilon}{2}}(0)$ orientation preserving.

We now claim that we may approximate $f_{\frac{\epsilon}{2}}$ arbitrarily well in C^k-norm on I_α^J by holomorphic automorphisms G of $\mathbb{C}^n$. Granted this for a moment, we proceed as follows. Let χ be a cutoff function with compact support in $\mathrm{int}(I_\alpha^J)$ such that $\chi \equiv 1$ near $Q_\alpha^J(\delta)$. Define

$$f'_{\frac{\epsilon}{2}}(x) := f_{\frac{\epsilon}{2}}(x) + \chi(x)(G(x) - f_{\frac{\epsilon}{2}}(x)). \tag{3.4}$$

We set $g_\alpha(z) := G_n(z)$, where G_n denotes the nth coordinate of G.

To produce G it suffices by [3] to produce a piecewise smooth isotopy G_t of embeddings connecting the identity map with f_α^J, such that $G_t(I_\alpha^J)$ is totally real and polynomially convex for all t. Let π_t be an isotopy of embeddings defined on $f_\alpha^J(I_\alpha^J)$ near the origin, such that π_0 is the identity map, and π_1 is the projection to $\mathbb{R}^d$. Define $\tilde{G}_t(x) := f_\alpha^J((1-t)x)$ for $t \in (0, 1-\eta)$ and compose $\tilde{G}_{1-\eta}$ with π_t. It η is sufficiently small, the whole isotopy is totally real and polynomially convex, and we have connected f_α^J to an orientation preserving embedding $H : I_\alpha^J \to \mathbb{R}^d$ with $H(0) = 0$. Now H is connected to $DH(0)$ via the path $\frac{1}{t}H(tx)$, and then $DH(0)$ is connected to the identity map.

Finally, note that there is plenty of room to perturb the automorphism G, e.g. by using shears, so we may assume that the intersection of any collection of $n+1$ zero sets is empty.

Second Step:

For simplicity we assume that M is without boundary. Let $I_1, ..., I_N$ be a cover of M by closed cubes, such that there is a collection of subcubes $I_i^0 \subset\subset \mathrm{int}(I_i)$ which also cover M. For each i let χ_i be a cutoff function compactly supported in $\mathrm{int}(I_i)$ with $\chi_i \equiv 1$ near I_i^0. Let $U_j \subset\subset U_{j+1} \subset U$ be open sets for $j = 1, ..., N$, and let Ω begiven by Proposition 3.2 for all the data $(f(M), U_{j+1}, U_j)$ with $j = 1, ..., N$. We will proceed to perturb the image $f(M)$ cube by cube.

Assume that we have constructed a small C^k-perturbation $f_j : M \to \mathbb{C}^n$ of f such that

$$h(K \cup f_j(\cup_{i=1}^j I_i^0)) \subset U_j$$

(it will be clear from the construction how to construct f_1).

By Step 1 we let $g_{j+1} : I_{j+1} \to \mathbb{C}^n$ be a small C^k-perturbation of f_j such that

$$h(K \cup f_j(\cup_{i=1}^j I_i^0) \cup g_{j+1}(I_{j+1})) \subset \Omega \cup U_j,$$

and g_{j+1} coincides with f_j in a neighbourhood of $\cup_{i=1}^j I_i^0$. Define f_{j+1} by

$$f_{j+1} = f_j + \chi_{j+1}(g_{j+1} - f_j).$$

4. Proof of Theorem 1.4

Let $(U_j)_{j\geq 1}$ be a sequence of open neighborhoods of K such that $U_{j+1} \subset U_j$ for all $j \geq 1$ and $K = \bigcap_{j\geq 1} U_j$. Set $f_0 := f$. We perturb f_0 inductively using Lemma 3.1, obtaining a family of totally real C^∞ embeddings $(f_j \colon M \to \mathbb{C}^n)_{j\in\mathbb{N}}$ such that,

(1) for all $j \geq 1$ we have $K \cup \widehat{f_j(M)} \subset U_j \cup f_j(M)$,
(2) for all $j \in \mathbb{N}$ we have
$$\|f_{j+1} - f_j\|_{C^{k+j}(M)} < \varepsilon_j,$$
where the sequence (ε_j) satisfies for all $\ell \in \mathbb{N}$,
$$\sum_{j=\ell}^{\infty} \varepsilon_j < \delta_\ell,$$
where $\delta_0 > 0$ is to be chosen and for all $\ell \geq 1$, δ_ℓ is the constant $c(f_\ell, K, U_\ell)$ given by Lemma 3.3,
(3) for all $j \in \mathbb{N}$ we have $f_{j+1} \equiv f_j$ on $f^{-1}(K)$.

The sequence (f_j) clearly converges to a C^∞ map $f' \colon M \to \mathbb{C}^n$ which coincides with f on $f^{-1}(K)$, and such that $\|f - f'\|_{C^k(M)} < \delta_0$. If δ_0 is small enough, then f' is a totally real C^∞ embedding. We claim that for all $\ell \geq 1$,
$$K \cup \widehat{f'(M)} \subset U_\ell \cup f'(M).$$
This follows from Lemma 3.3, since
$$\|f' - f_\ell\|_{C^1(M)} \leq \sum_{j=\ell}^{\infty} \varepsilon_j < \delta_\ell.$$

□

References

1. M. Adachi, *Embeddings and Immersions*, Translations of Mathematical Monographs **124**
2. H. Alexander, *Hulls of subsets of the torus in* $\mathbb{C}^2$, Ann. Inst. Fourier (Grenoble) **48** (1998), no. 3, 785–795.
3. F. Forstneric, *Approximation by automorphisms on smooth submanifolds of* $\mathbb{C}^n$, Math. Ann. 300 (1994), 719–738.
4. F. Forstneric, J.-P. Rosay, *Approximation of biholomorphic mappings by automorphisms of* $\mathbb{C}^n$, Invent. Math. **112** (1993), no. 2, 323–349.
5. P. Gupta, *A nonpolynomially convex isotropic two-torus with no attached discs*, arXiv:1610.09356.
6. P. T. Ho, H. Jacobowitz, P. Landweber, *Optimality for totally real immersions and independent mappings of manifolds into* $\mathbb{C}^N$, New York J. Math. **18** (2012) 463–477.
7. A. Izzo, H. Samuelsson Kalm, E. F. Wold, *Presence or absence of analytic structure in maximal ideal spaces*, Math. Ann. **366** (2016), no. 1-2, 459–478.
8. A. Izzo, E. L. Stout, *Hulls of surfaces*, Indiana U. Math. J. (to appear).
9. E. Løw, E. F. Wold, *Polynomial convexity and totally real manifolds*, Complex Var. Elliptic **54** (2009), Nos. 3–4, 265–281.
10. H. Whitney, *Differentiable manifolds*, Ann. of Math. (2) **37** (1936), no. 3, 645–680.

L. Arosio: Dipartimento Di Matematica, Università di Roma "Tor Vergata", Via Della Ricerca Scientifica 1, 00133, Roma, Italy
E-mail address: arosio@mat.uniroma2.it

E. F. Wold: Department of Mathematics, University of Oslo, Postboks 1053 Blindern, NO-0316 Oslo, Norway
E-mail address: erlendfw@math.uio.no